

## RECOMMENDED PREVENTIVE SEXUAL HEALTH SERVICES FOR WOMEN

Sexual Health Service	Adolescent (13-18 years)	Adult (19+)
<b>Cervical cancer screening</b>	Not recommended	Those age 21-29 every 3 years Those age 30-65 every 3-5 years
<b>Chlamydia screening</b>	Those who are sexually active (screen annually) or pregnant	Those aged 25 and under (screen annually) Those over age 26 only if they are at risk (screen annually) Those who are pregnant
<b>Contraceptive counseling</b>	Those who are sexually active	Those who are sexually active and/or of childbearing age
<b>Counseling to prevent sexually transmitted infections</b>	Those who are sexually active	Those who are at risk
<b>Gonorrhea screening</b>	Those who are sexually active or pregnant	Those who are at risk, including those who are pregnant
<b>Hepatitis A vaccine</b>	Those not vaccinated as infants	Those who are at risk
<b>Hepatitis B vaccine</b>	Those not vaccinated as infants	Those who are at risk Those who are unvaccinated and have diabetes
<b>Hepatitis B screening</b>	Those who are pregnant	Those who are pregnant
<b>Hepatitis C screening</b>	Not recommended	Those born between 1945 and 1965
<b>HIV testing</b>	Those who are or have been sexually active (test at least once) or are pregnant	Those who are or have been sexually active (test at least once) Those who are at risk (test at least annually) Those who are pregnant
<b>HPV vaccine</b>	Girls aged 11-12 and unvaccinated teens	May be given through age 26 (and to eligible adults over age 26)
<b>Syphilis screening</b>	Those who are at risk or pregnant	Those who are at risk or pregnant

You're at risk if you have had unprotected vaginal, anal, or oral sex (sex without a condom), have multiple partners, have an STI (including HIV), shared needles, exchanged sex for drugs or money, or have a partner who answers "yes" to any of these questions or whose health status you don't know.