RECOMMENDED PREVENTIVE SEXUAL HEALTH SERVICES FOR MEN

| Sexual Health Service | Adolescent (13-18 years) | Adult (19+) |
|---|---|--|
| Chlamydia screening | Those who are at risk | Those who are at risk |
| Counseling to prevent sexually transmitted infections | Those who are sexually active | Those who are at risk |
| Gonorrhea screening | Those who are at risk | Those who are at risk |
| Hepatitis A vaccine | Those not vaccinated as infants | Men who have sex with men |
| Hepatitis B vaccine | Those not vaccinated as infants | Those who are at risk and are unvaccinated Those who are unvaccinated and have diabetes |
| Hepatitis B screening | Not recommended | Men who have sex with men and are unvaccinated |
| Hepatitis C screening | Not recommended | Those born between 1945 and 1965 Men who have sex with men, especially those that have used injection drugs or are HIV+ |
| HIV testing | Those who have been or are sexually active (test at least once) | Those who have been or are sexually active (test at least once) |
| HPV vaccine | Boys aged 11-12 and unvaccinated teens | May be given through age 26 |
| Syphilis screening | Those who are at risk | Those who are at risk |

How Do I Know if I'm at Risk?

Many behaviors put you at risk for getting STIs, including HIV. You are at risk if you answer "yes" to **any** of the following questions:

- Have you had unprotected vaginal, anal, or oral sex (sex without a condom)?
- Do you have multiple partners?
- Do you have an STI, including HIV?

- Have you shared needles?
- Do you exchange sex for drugs or money?
- Do you have a partner who answers "yes" to any of these questions or whose health status you don't know?